


# Two-Photon Physics at



Richard Nisius, CERN  
Aachen, 28 Januar 1997

## 1. Photon-Photon scattering

- Exclusive hadronic final states
- Inclusive hadronic final states

## 2. Electron-Photon DIS

- Lepton pairs and  $F_{2,QED}^\gamma$
- The structure function  $F_2^\gamma(x, Q^2)$

# Analysis topics in Two-Photon events at LEP


	$\gamma\gamma$ scattering		$e\gamma$ scattering	
	exclusive hadronic f.s.	lepton pairs $\gamma\gamma \rightarrow$ hadrons	lepton pairs $\gamma^* \gamma \rightarrow$ hadrons	singly tagged events
A	$D^*(2010)^\pm$	hadron flow		
D		hadron flow	$\mu$ $F_{2,QED}^\gamma$	$F_2^\gamma$ hadron flow
L	$K_S^0 K_S^0, \eta'(958)$ $a_2(1320), f_2(1720)?$ $\eta_c(1S), \chi_{c2}(1P)$	$e\mu\tau$ $\sigma(W_{\gamma\gamma})$	$e, \mu$ $F_{2,QED}^\gamma$	
O		$\frac{d\sigma}{d\eta^{jet}} \frac{d\sigma}{dE_T^{jet}}$	$e\mu\tau$ $F_{2,QED}^\gamma$	$F_2^\gamma$ hadron flow

preliminary

published

presented

## Photon–photon scattering



Exchange of two quasi-real photons ( $\gamma$ )


$$Q_i^2 = 2E_i E'_i (1 - \cos \theta_i) \approx 0$$

$$W^2 = s_{\gamma\gamma} = \left( \sum_h E_h \right)^2 - \left( \sum_h \vec{p}_h \right)^2$$

At  $\sqrt{s_{ee}} = 130 \text{ GeV}$ , for  $W^2 > 4 \text{ GeV}^2$  and  $Q_i^2 < 1 \text{ GeV}^2$ :

$$\sigma(e^+ e^- \rightarrow e^+ e^- + \text{hadrons}) \approx 14 \text{ nb} \approx 40 \cdot \sigma(e^+ e^- \rightarrow (\gamma, Z^0) \rightarrow \text{hadrons})$$

# $D^*(2010)^\pm$ production at LEP1



slow pion

$$D^{*+} \rightarrow D^0 \pi^+$$



Decaymodes:

(1)  $D^0 \rightarrow K^- \pi^+$

(2)  $D^0 \rightarrow K^- \pi^+ \pi^0$

(3)  $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$



$$M(D^{*\pm}) - M(D^0)$$



$$\sigma(e^+e^- \rightarrow e^+e^- D^{*\pm} X) = 155 \pm 33 \pm 21 \text{ pb}$$

$$\Rightarrow \sigma(e^+e^- \rightarrow e^+e^- c\bar{c}X) = 326 \pm 87 \text{ pb}$$

# $K_S^0 K_S^0$ final states



$$\sigma(\gamma^* \gamma^* \rightarrow R) = 8\pi(2J_R + 1) \frac{\Gamma_{\gamma\gamma}(R)\Gamma(R)}{(W_{\gamma\gamma}^2 - m_R^2)^2 + m_R^2 \Gamma^2(R)}$$


$$f'_2(1525) \rightarrow K_S^0 K_S^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

$$f'_2(1525) \text{ signal of } 31 \pm 6 \text{ events for } \mathcal{L}_{int} = 114 \text{ pb}^{-1}$$


$$\Gamma_{\gamma\gamma}(f'_2) \cdot BR(f'_2 \rightarrow K \bar{K}) = (0.093 \pm 0.018 \pm 0.022) \text{ keV}$$

Probably pure helicity 2 state

# $\gamma\gamma \rightarrow \text{hadrons at } \sqrt{s_{ee}} = M_{Z^0}$



## DELPHI




# Leading order diagrams


Direct:



Single-Resolved:



Double-Resolved:



## **Monte Carlo models**

### **PYTHIA 5.721 and PHOJET 1.05**

#### Monte Carlo ingredients:

1. Leading order (LO) QCD matrix elements
2. Hard and soft processes
3. Total cross sections from Regge models
4. Initial state parton radiation
5. Fragmentation based on by JETSET 7.408
6. Multiple interactions


## **NLO calculations**

- NLO calculations for inclusive jet cross sections by T. Kleinwort and G. Kramer, DESY-96-035 (1996), hep-ph/9509321 and Phys. Lett. B370 (1996) 141, hep-ph/9602418.

# The $x_\gamma$ distribution for 2-jet events at $\sqrt{s_{ee}} = 131 \text{ GeV}$

$x_\gamma$  is the fraction of the photon momentum participating in the hard interaction

$$x_\gamma = \max \left\{ \frac{\sum_{\text{jets}} (E + p_z)}{\sum_{\text{hadrons}} (E + p_z)}, \frac{\sum_{\text{jets}} (E - p_z)}{\sum_{\text{hadrons}} (E - p_z)} \right\}$$




**Direct events :**  $x_\gamma \equiv 1$  no remnant jet

**Resolved events :**  $x_\gamma < 1$  remnant jets possible

# The separation of event classes


at  $\sqrt{s_{ee}} = 131 \text{ GeV}$



$$x_{\gamma}^{\pm} = \frac{\sum_{\text{jets}} (E \pm p_z)}{\sum_{\text{hadrons}} (E \pm p_z)}$$



# A direct two-jet event

Run: event 6839: 71842 Date 951109 Time 135804 Ctrk(N= 10 Sump= 9.8) Ecal(N= 25 SurfE= 15.1) Hcal(N= 6 SurfE= 2.3)  
 Ebeam 65.129 Evis 23.2 Emiss 107.0 Vtx ( -.03, .08, -.59) Muon(N= 0) Sec Vtx(N= 0) Fdet(N= 0 SurfE= .0)  
 Bz=4.350 Bunchlet 1/1 Thrust= .7091 Aplan= .0339 Oblat= .5027 Spher= .7239




x-y view

Side view - plane of Thrust axis



Side view - plane perp. to Thrust

- CTRK OPAL Run 6839 Event 71842 -
- ECAL
- HCAL
- FDET+SiW



**Jet 1 :**

$$\eta = 0.9, E_T^{\text{jet}} = 6.6 \text{ GeV}$$


**Jet 2 :**

$$\eta = 0.7, E_T^{\text{jet}} = 6.9 \text{ GeV}$$


# The inclusive jet cross-sections

at  $\sqrt{s_{ee}} = 133 \text{ GeV}$

$\frac{d\sigma}{d\eta^{\text{jet}}}$  compared to Monte Carlo models




$\frac{d\sigma}{dE_T^{\text{jet}}}$  compared to NLO Calculations




# The inclusive 2-jet cross-sections

at  $\sqrt{s_{ee}} = 161 \text{ GeV}$


$\frac{d\sigma}{d\eta^{\text{jet}}}$  compared to Monte Carlo models



$\frac{d\sigma}{dE_T^{\text{jet}}}$  compared to NLO Calculations



# Electron-Photon Scattering



$$\frac{d^2 \sigma_{e\gamma \rightarrow eX}}{dx dQ^2} = \frac{2\pi\alpha^2}{x Q^4} \cdot$$


$$\left[ (1 + (1 - y)^2) F_2^\gamma(x, Q^2) - \underbrace{y^2 F_L^\gamma(x, Q^2)}_{\rightarrow 0} \right]$$

$$Q^2 = 2 E_b E_{\text{tag}} (1 - \cos \theta_{\text{tag}}) \gg P^2$$

$$x = \frac{Q^2}{Q^2 + W^2 + P^2}$$

$$y = 1 - \frac{E_{\text{tag}}}{E_b} \cos^2\left(\frac{\theta_{\text{tag}}}{2}\right) \ll 1$$

## The production of lepton pairs








$$\frac{d^2\sigma_{e\gamma \rightarrow e\mu^+\mu^-}}{dx dQ^2} = \frac{2\pi\alpha^2}{x Q^4} \left[ (1 + (1 - y)^2) F_{2,\text{QED}}^\gamma - y^2 F_{L,\text{QED}}^\gamma \right]$$


$$F_{2,\text{QED}}^\gamma(x, Q^2, P^2 = 0)/\alpha \approx \frac{x}{\pi} \left[ 1 - 2x(1 - x) \ln \frac{W^2}{m_\mu^2} - 1 + 8x(1 - x) \right]$$

$$F_{L,\text{QED}}^\gamma(x, Q^2, P^2 = 0)/\alpha \approx \frac{4}{\pi} x^2(1 - x)$$

# The world data on $F_2^{\gamma}$ <sub>QED</sub> compared to QED



# The structure function for $P^2 > 0$




$\theta_{\text{antitag}} = 0.026$  rad

$P^2 (0.25E_b) = 1.25 \text{ GeV}^2$

$P^2 (0.25E_b) = 0.35 \text{ GeV}^2$


$P^2 (0.23E_b) = 0.05 \text{ GeV}^2$




$F_{2,\text{QED}}^\gamma(x, Q^2, P^2)/\alpha \approx$

$$\frac{x}{\pi} \left[ 1 - 2x(1-x) \ln \frac{W^2}{m_\mu^2 + x(1-x)P^2} - 1 + 8x(1-x) - \frac{x(1-x)P^2}{m_\mu^2 + x(1-x)P^2} \right]$$

# The contribution of $F_{L,QED}^\gamma$





# Azimuthal Correlations




**OPAL**

arbitrary scales



$\chi$

# The measurement of $F_B^\gamma / F_2^\gamma$




$$\frac{d\sigma(e\gamma \rightarrow e\mu^+\mu^-)}{dx dy d\chi/2\pi} \simeq \frac{2\pi\alpha^2}{Q^2} \left( \frac{1 + (1-y)^2}{xy} \right) \cdot F_2^\gamma \left( 1 + \frac{1}{2}\epsilon(F_B^\gamma/F_2^\gamma) \cos 2\chi \right)$$


$$\frac{1}{2}\epsilon = \frac{(1-y)}{1+(1-y)^2} \approx 1, \quad F_B^\gamma = F_L^\gamma = \frac{4\alpha}{\pi} x^2 (1-x) \quad (\text{in LO})$$

# The contributions to $F_2^\gamma(x, Q^2)$

$$F_2^\gamma(x, Q^2) = x \sum_{c,f} e_q^2 \underline{f_{q,\gamma}(x, Q^2)}$$



hadronic, VDM,  $p_T = \text{"small"}$  pointlike,  $p_T = \text{"large"}$   
 $\rho, \omega, \phi$ , non-perturbative perturbative



## Event selection

$$\mathcal{L}_{int} = 156.4 \text{ pb}^{-1}$$

1. Electron Tag:  $E_{\text{tag}} \geq 0.775 E_b$  and  
 $0.06 \leq \theta_{\text{tag}} \leq 0.12 \text{ rad}$

2. Antitag:  $E_a \leq 0.25 E_b$

3.  $N_{\text{ch}} \geq 3$ , and  $(2.5 \leq W_{\text{vis}} \leq 40) \text{ GeV}$

$\Rightarrow 5455 \text{ events with } (6 \lesssim Q^2 \lesssim 30) \text{ GeV}^2$

1. Electron Tag:  $0.75 E_b \leq E_{\text{tag}} \leq 1.15 E_b$  and  
 $0.2 \leq \theta_{\text{tag}} \leq 0.5 \text{ rad}$ , plus isolation criteria


2. Antitag:  $E_a \leq 0.15 E_b$

3.  $N_{\text{ch}} \geq 3$ , and  $(2.5 \leq W_{\text{vis}} \leq 25) \text{ GeV}$


4.  $p_{\text{t,bal}} < 5 \text{ GeV}$ ,  $p_{\text{t,out}} < 4 \text{ GeV}$ ,  
 $-0.5 E_b \leq p_{\text{z,miss}} \leq 0.5 E_b$

$\Rightarrow 225 \text{ events with } (60 \lesssim Q^2 \lesssim 400) \text{ GeV}^2$

# The phase space at $\sqrt{s_{ee}} = M_{Z^0}$



# The effect of different $E_\gamma$




# A tagged two-photon event

Run: event 6422: 47694 Date 950817 Time 155240 Ctrk(N= 8 Surp= 12.4) Ecal(N= 19 SurE= 47.2) Hcal(N= 6 SurE= 3.4)  
 Ebeam 45.595 Evis 58.2 Emiss 33.0 Vtx ( -0.05, 0.11, 1.17) Muon(N= 0) Sec Vtx(N= 0) Fdet(N= 0 SurE= 0.0)  
 Bz=4.028 Bunchlet 1/1 Thrust=0.7818 Aplan=0.0006 Oblat=0.4802 Spher=0.0371






## Event type bits

- 4: Low mult. preselect
- 8: Singl phot preselect
- 12: Tagged two phot
- 13: Higgs high mult
- 24: S phot EM ass TOF
- 25: S phot EM and TOF
- 26: S phot In-time TOF
- 27: S phot EM elus
- 28: S phot High pT trk
- 30: S phot no HMMJ vet
- 31: long-lived decays
- 32: "Phys1" selection
- 1: Z0 type physics



Status  
 Det Tr  
 CTRK  
 ECAL  
 HCAL  
 FDET+SiW



## Kinematics :





$$E_{tag} = 39.2 \text{ GeV}$$

$$\theta_{tag} = 21.3^\circ$$


$$Q^2 = 245 \text{ GeV}^2$$

$$W_{vis}^2 = 17 \text{ GeV}^2$$

# Event distributions compared to HERWIG





# The flow of hadronic energy as predicted by HERWIG




Only about 10% of the energy is deposited outside of  
the detector acceptance

# The $W - W_{\text{vis}}$ correlation




The correlation based on F2GEN is much stronger  
The inclusion of the **Forward Region** significantly  
improves the correlation

# The energy flow for $\sqrt{s_{ee}} = M_{Z^0}$





# The model dependence as a function of $x_{vis}$



The agreement gets better for increasing  $x_{vis}$

# The corrected flow of hadronic energy



The description of the data by the Monte Carlo models is poor at low  $x$  and  $Q^2$  and it improves for higher  $Q^2$ . The data, however, is precise enough to further constrain the models!

## Some words about unfolding

### The Principle:

$$g^{\text{det}}(\mathbf{u}) = \int A(\mathbf{u}, \omega) f^{\text{part}}(\omega) d\omega + B(\mathbf{u})$$

#### 1. Our case:

$g^{\text{det}}(\mathbf{u}) = g^{\text{det}}(\mathbf{x}_{\text{vis}})$ ,  $\mathbf{x}_{\text{vis}} = f(E_{\text{tag}}, \theta_{\text{tag}}, W_{\text{vis}})$   
and  $f^{\text{part}}(\omega) = f^{\text{part}}(\mathbf{x})$  which is related to  $F_2^\gamma$ ,  $B(\mathbf{u})$   
is denotes the background events.


2.  $A(\mathbf{u}, \omega)$  has to be obtained from the Monte Carlo Models  
 $\Rightarrow$  **Model Dependence**, consider all reasonable models.

3. The  $g^{\text{det}}(\mathbf{x}_{\text{vis}})$  distribution from the Monte Carlo is  
changed during unfolding, by assigning weights to each  
Monte Carlo event, in order to match the  $g^{\text{det}}(\mathbf{x}_{\text{vis}})$   
distribution of the data.

- The  **$g^{\text{det}}(\mathbf{x}_{\text{vis}})$  distributions** of data and Monte Carlo  
**agree** afterwards by construction.
- **Other distributions** have to be used in order to check  
whether the unfolding has also improved on them,  
**without** using explicitly this variable.


4. The unfolding result **should** be independent of the  $F_2^\gamma$  used  
in the Monte Carlo. This is **not** true if  $F_2^\gamma$  and the  $\gamma^* \gamma$   
fragmentation do **not factorize**.

# The $x_{\text{vis}}$ distributions compared to HERWIG



The mean  $x_{\text{vis}}$  increases with  $Q^2$   
The background contributions are small


# Unfolding tests using HERWIG with GRV as unfolding MC



The error is dominated by systematic effects

# OPAL results on $F_2^\gamma(x, Q^2)$







at  $\sqrt{s_{ee}} = M_{Z^0}$



$$F_2^\gamma(Q^2)/\alpha = (0.08^{+0.13}_{-0.18}) + (0.13^{+0.06}_{-0.04}) \ln Q^2$$

$\chi^2/\text{dof} = 0.05$   $Corr = -0.95$

# A selection of $F_2^\gamma(x, Q^2)$ measurements compared to GRV (LO)




## The systematic error on $F_2^\gamma(x, Q^2)$

$\langle Q^2 \rangle$ (GeV <sup>2</sup> )	$\langle x \rangle$	$x$ - range	$F_2^\gamma / \alpha$	rel (%)
<b>PLUTO</b> 9.2	0.145	0.060-0.230	$0.35 \pm 0.03 \pm 0.09$	25
	0.385	0.230-0.540	$0.40 \pm 0.03 \pm 0.06$	15
	0.720	0.540-0.900	$0.49 \pm 0.07 \pm 0.07$	15
<b>OPAL</b> 14.7	0.072	0.006-0.137	$0.38 \pm 0.01 \begin{smallmatrix} +0.06 \\ -0.13 \end{smallmatrix}$	25
	0.230	0.137-0.324	$0.41 \pm 0.02 \begin{smallmatrix} +0.06 \\ -0.03 \end{smallmatrix}$	11
	0.423	0.324-0.522	$0.41 \pm 0.03 \begin{smallmatrix} +0.08 \\ -0.11 \end{smallmatrix}$	23
	0.679	0.522-0.836	$0.54 \pm 0.05 \begin{smallmatrix} +0.31 \\ -0.13 \end{smallmatrix}$	41
<b>TOPAZ</b> 16	0.085	0.020-0.150	$0.60 \pm 0.08 \pm 0.06$	10
	0.240	0.150-0.330	$0.56 \pm 0.09 \pm 0.04$	7
	0.555	0.330-0.780	$0.46 \pm 0.15 \pm 0.06$	13


The single contributions for OPAL  $\langle Q^2 \rangle = 14.7 \text{ GeV}^2$

$\langle x \rangle$	0.072	0.230	0.423	0.679
$E_{\text{tag}}$	$+0.02$ $-0.01$	$+0.01$ $-0.02$	$+0.02$ $-0.02$	$+0.02$ $-0.01$
$\theta_{\text{tag}}$	$< 0.01$	$-0.02$	$+0.04$	$< +0.01$ $-0.10$
$W_{\text{vis}}^2$	$< 0.01$	$-0.02$	$+0.05$	$-0.12$
$p_t$	$< 0.01$	$< 0.01$	$< 0.01$	$< 0.01$
PDF	$+0.06$	$+0.02$	$-0.11$	$-0.13$
SUE	$-0.05$	$-0.03$	$< 0.01$	$+0.02$
model	$+0.03$ $-0.13$	$+0.06$	$+0.08$ $-0.03$	$+0.31$

# $dF_2^\gamma / d \ln Q^2$ for $n_f = 4 = (udsc)$




# $dF_2^\gamma / d \ln Q^2$ for $n_f = 3 = (uds)$





# The prospects of $F_2^\gamma$ at higher $E_b$



## Conclusions

Two-Photon physics is a very active field at LEP with good prospects for LEP2

- Photon-Photon scattering

1. A number of resonances have been measured.
2. The flow of hadronic energy has been compared to Monte Carlo models.
3. NLO calculations of jet production agree nicely with the data.

- Electron-Photon DIS

1. There is in good agreement with QED predictions and the measured  $F_{2,QED}^\gamma$  structure function and the ratio  $F_B^\gamma / F_2^\gamma$ .
2. The  $F_2^\gamma$  structure function was measured for  $7.5 < \langle Q^2 \rangle < 135 \text{ GeV}^2$ . The systematic errors have a large contribution from the imperfect description of the hadronic final state by the QCD inspired Monte Carlo models.

and . . .

# Outlook

## What can we expect from LEP on Two-Photon physics in the future

- Photon-Photon scattering

1. More resonances (see list).
2. Jet production for the direct component alone.
3. Determination of the gluon content of the photon in jet production.
4. . . .

- Electron-Photon DIS

1.  $P^2$  dependence of  $F_{2,\text{QED}}^\gamma$ .
2. Azimuthal correlations in hadronic final states.
3.  $F_2^\gamma$  for  $20 < Q^2 < 1000 \text{ GeV}^2$ .
4. Double tag events.
5. . . .

**The LEP2 programme has just started**

slides:

<http://wwwcn1.cern.ch/~nisius/talks/AACH280197/index.html>